



## EIF2AK4 gene

eukaryotic translation initiation factor 2 alpha kinase 4

### Normal Function

The *EIF2AK4* gene provides instructions for making a protein that helps direct a cell's response to changes that could damage the cell. This protein is found in several tissues throughout the body, including blood vessel walls. The EIF2AK4 protein can turn on (activate) another protein called eIF2 alpha (eIF2 $\alpha$ ), which helps control protein production. When cells are under stress, for example when the level of protein building blocks (amino acids) is too low, EIF2AK4 activates eIF2 $\alpha$ . When turned on, eIF2 $\alpha$  stimulates processes that reduce protein production, which helps conserve amino acids. In addition, activated eIF2 $\alpha$  can trigger production of certain proteins called transcription factors, which control gene activity. The transcription factors regulated by eIF2 $\alpha$  control the activity of genes involved in processes that help reduce the stress on the cell.

### Health Conditions Related to Genetic Changes

pulmonary arterial hypertension

pulmonary veno-occlusive disease


Mutations in the *EIF2AK4* gene are the primary genetic cause of a condition called pulmonary veno-occlusive disease (PVOD); at least 22 mutations in this gene have been found in affected individuals. In PVOD, excess fibrous tissue builds up in the small vessels in the lungs that carry oxygenated blood from the lungs to the heart (the pulmonary veins). This buildup narrows (occludes) the vessels and impairs blood flow. Because blood flow through the lungs is difficult, pressure rises in the vessels that carry blood that needs to be oxygenated to the lungs from the heart (pulmonary arteries). Increased pressure in these vessels is known as pulmonary arterial hypertension.

The *EIF2AK4* gene mutations involved in PVOD likely lead to a complete loss of functional protein. It is not known how absence of EIF2AK4 protein function leads to the pulmonary vein abnormalities characteristic of PVOD.

## Chromosomal Location

Cytogenetic Location: 15q15.1, which is the long (q) arm of chromosome 15 at position 15.1

Molecular Location: base pairs 39,934,101 to 40,035,596 on chromosome 15 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- E2AK4\_HUMAN
- eukaryotic translation initiation factor 2-alpha kinase 4
- GCN2
- GCN2 eIF2alpha kinase
- GCN2-like protein
- general control nonderepressible 2
- KIAA1338
- PVOD2

## Additional Information & Resources

### Educational Resources

- Madame Curie Biosciences Database (2000): Mechanism of Translation Initiation in Eukaryotes  
<https://www.ncbi.nlm.nih.gov/books/NBK6597/>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28EIF2AK4%5BTIAB%5D%29+OR+%28eukaryotic+translation+initiation+factor+2+alpha+kinase+4%5BTIAB%5D%29%29+OR+%28%28GCN2+eIF2alpha+kinase%5BTIAB%5D%29+OR+%28GCN2%5BTIAB%5D%29+OR+%28eukaryotic+translation+initiation+factor+2+alpha+kinase+4%5BTIAB%5D%29+OR+%28general+control+nonderepressible+2%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

### OMIM

- EUKARYOTIC TRANSLATION INITIATION FACTOR 2-ALPHA KINASE 4  
<http://omim.org/entry/609280>

### Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC EIF2AK4.html](http://atlasgeneticsoncology.org/Genes/GC	EIF2AK4.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=EIF2AK4%5Bgene%5D>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=19687](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=19687)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/440275>
- UniProt  
<http://www.uniprot.org/uniprot/Q9P2K8>

### **Sources for This Summary**

- Deval C, Chaveroux C, Maurin AC, Cherasse Y, Parry L, Carraro V, Milenkovic D, Ferrara M, Bruhat A, Jousse C, Fafournoux P. Amino acid limitation regulates the expression of genes involved in several specific biological processes through GCN2-dependent and GCN2-independent pathways. *FEBS J.* 2009 Feb;276(3):707-18. doi: 10.1111/j.1742-4658.2008.06818.x. Epub 2008 Dec 19.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/19120448>
- Donnelly N, Gorman AM, Gupta S, Samali A. The eIF2 $\alpha$  kinases: their structures and functions. *Cell Mol Life Sci.* 2013 Oct;70(19):3493-511. doi: 10.1007/s00018-012-1252-6. Epub 2013 Jan 26. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23354059>
- OMIM: EUKARYOTIC TRANSLATION INITIATION FACTOR 2-ALPHA KINASE 4  
<http://omim.org/entry/609280>

- Eyries M, Montani D, Girerd B, Perret C, Leroy A, Lonjou C, Chelghoum N, Coulet F, Bonnet D, Dorfmüller P, Fadel E, Sitbon O, Simonneau G, Tregouët DA, Humbert M, Soubrier F. EIF2AK4 mutations cause pulmonary veno-occlusive disease, a recessive form of pulmonary hypertension. *Nat Genet.* 2014 Jan;46(1):65-9. doi: 10.1038/ng.2844. Epub 2013 Dec 1.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/24292273>
  - Murguía JR, Serrano R. New functions of protein kinase Gcn2 in yeast and mammals. *IUBMB Life.* 2012 Dec;64(12):971-4. doi: 10.1002/iub.1090. Epub 2012 Nov 5. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23129244>
- 

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/EIF2AK4>

Reviewed: March 2015

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services